

8. Kiyoi H, Yanada M, Ozekia K. Clinical significance of FLT3 in leukemia. *Int J Hematol*. 2005;82:85-92.
9. Yanada M, Matsuo K, Suzuki T, Kiyoi H, Naoe T. Prognostic significance of FLT3 internal tandem duplication and tyrosine kinase domain mutations for acute myeloid leukemia: a meta-analysis. *Leukemia*. 2005;19:1345-1349.
10. Frohling S, Schlenk RF, Breitnick J, et al. Prognostic significance of activating FLT3 mutations in younger adults (16 to 60 years) with acute myeloid leukemia and normal cytogenetics: a study of the AML Study Group Ulm. *Blood*. 2002;100:4372-4380.
11. Murphy KM, Levis M, Hafez MJ, et al. Detection of FLT3 internal tandem duplication and D835 mutations by a multiplex polymerase chain reaction and capillary electrophoresis assay. *J Mol Diagn*. 2003;5:96-102.
12. Slovak ML, Kopecky KJ, Cassileth PA, et al. Karyotypic analysis predicts outcome of preremission and postremission therapy in adult acute myeloid leukemia: a Southwest Oncology Group/Eastern Cooperative Oncology Group Study. *Blood*. 2000;96:4075-4083.
13. Karp JE, Blackford A, Smith BD, et al. Clinical activity of sequential flavopiridol, cytosine arabinoside, and mitoxantrone for adults with newly diagnosed, poor-risk acute myelogenous leukemia. *Leuk Res*. 2010;34:877-882.
14. Vaughan WP, Karp JE, Burke PJ. Two-cycle timed-sequential chemotherapy for adult acute nonlymphocytic leukemia. *Blood*. 1984;64:975-980.
15. Castaigne S, Chevret S, Archimbaud E, et al. Randomized comparison of double induction and timed-sequential induction to a "3 + 7" induction in adults with AML: long-term analysis of the Acute Leukemia French Association (ALFA) 9000 study. *Blood*. 2004;104:2467-2474.
16. McCune JS, Batchelder A, Deeg HJ, et al. Cyclophosphamide following targeted oral busulfan as conditioning for hematopoietic cell transplantation: pharmacokinetics, liver toxicity, and mortality. *Biol Blood Marrow Transplant*. 2007;13:853-862.
17. Luznik L, O'Donnell PV, Symons HJ, et al. HLA-haploidentical bone marrow transplantation for hematologic malignancies using nonmyeloablative conditioning and high-dose, posttransplantation cyclophosphamide. *Biol Blood Marrow Transplant*. 2008;14:641-650.
18. Luznik L, Fuchs EJ. High-dose, post-transplantation cyclophosphamide to promote graft-host tolerance after allogeneic hematopoietic stem cell transplantation. *Immunol Res*. 2010;47:65-77.
19. Gale RE, Hills R, Kottaridis PD, et al. No evidence that FLT3 status should be considered as an indicator for transplantation in acute myeloid leukemia (AML): an analysis of 1135 patients, excluding acute promyelocytic leukemia, from the UK MRC AML10 and 12 trials. *Blood*. 2005;106:3658-3665.
20. Meshinchi S, Arceci RJ, Sanders JE, et al. Role of allogeneic stem cell transplantation in FLT3/ITD-positive AML. *Blood*. 2006;108:400-401.
21. Bornhauser M, Illmer T, Schaich M, et al. Improved outcome after stem-cell transplantation in FLT3/ITD-positive AML. *Blood*. 2007;109:2264-2265.
22. Fernandez HF, Sun Z, Yao X, et al. Anthracycline dose intensification in acute myeloid leukemia. *N Engl J Med*. 2009;361:1249-1259.
23. Santos FP, Jones D, Qiao W, et al. Prognostic value of FLT3 mutations among different cytogenetic subgroups in acute myeloid leukemia. *Cancer*. 2010.
24. Gaidzik V, Dohner K. Prognostic implications of gene mutations in acute myeloid leukemia with normal cytogenetics. *Semin Oncol*. 2008;35:346-355.
25. Mead AJ, Linch DC, Hills RK, et al. FLT3 tyrosine kinase domain mutations are biologically distinct from and have a significantly more favorable prognosis than FLT3 internal tandem duplications in patients with acute myeloid leukemia. *Blood*. 2007;110:1262-1270.

## Predictions in the Face of Clinical Reality: *HistoCheck* versus High-Risk HLA Allele Mismatch Combinations Responsible for Severe Acute Graft-versus-Host Disease

Medhat Askar,<sup>1</sup> Ronald Sobecks,<sup>2</sup> Yasuo Morishima,<sup>3</sup> Takakazu Kawase,<sup>3</sup>  
Amy Nowacki,<sup>4</sup> Hideki Makishima,<sup>5</sup> Jaroslaw Maciejewski<sup>5</sup>

HLA polymorphism remains a major hurdle for hematopoietic stem cell transplantation (HSCT). In 2004, Elsner et al. proposed the *HistoCheck* Web-based tool to estimate the allogeneic potential between HLA-mismatched stem cell donor/recipient pairs expressed as a sequence similarity matching (SSM). SSM is based on the structure of HLA molecules and the functional similarity of amino acids. According to this algorithm, a high SSM score represents high dissimilarity between MHC molecules, resulting in a potentially more

From the <sup>1</sup>Transplant Center, Cleveland Clinic, Cleveland, Ohio; <sup>2</sup>Taussig Cancer Institute, Cleveland Clinic, Cleveland, Ohio; <sup>3</sup>Aichi Cancer Center, Nagoya, Japan; <sup>4</sup>Department of Quantitative Health Sciences, Cleveland Clinic, Cleveland, Ohio; and <sup>5</sup>Department of Translational Hematology and Oncology Research, Cleveland Clinic, Cleveland, Ohio.

**Financial disclosure:** See Acknowledgments on page 1414.

Correspondence and reprint requests: Medhat Askar, MD, PhD, D(ABHI), D(ABMLI), HCLD(ABB), Allogen Laboratories, Transplant Center, Cleveland Clinic, Department of Surgery, Cleveland Clinic Lerner College of Medicine, Case Western Reserve University, 9500 Euclid Ave, C100 MZA, Cleveland, OH 44195 (e-mail: [AskarM@CCF.ORG](mailto:AskarM@CCF.ORG)).

Received August 31, 2010; accepted January 26, 2011

© 2011 American Society for Blood and Marrow Transplantation

1083-8791/\$36.00

doi:10.1016/j.bbmt.2011.01.015

deleterious impact on stem cell transplant outcomes. We investigated the potential of SSM to predict high-risk HLA allele mismatch combinations responsible for severe acute graft-versus-host disease (aGVHD grades III and IV) published by Kawase et al., by comparing SSM in low- and high-risk combinations. SSM was calculated for allele mismatch combinations using the *HistoCheck* tool available on the Web ([www.histocheck.org](http://www.histocheck.org)). We compared ranges and means of SSM among high-risk (15 combinations observed in 722 donor/recipient pairs) versus low-risk allele combinations (94 combinations in 3490 pairs). Simulation scenarios were created where the recipient's HLA allele was involved in multiple allele mismatch combinations with at least 1 high-risk and 1 low-risk mismatch combination. SSM values were then compared. The mean SSM for high- versus low-risk combinations were 2.39 and 2.90 at A, 1.06 and 2.53 at B, 16.60 and 14.99 at C, 4.02 and 3.81 at DRB1, and 7.47 and 6.94 at DPB1 loci, respectively. In simulation scenarios, no predictable SSM association with high- or low-risk combinations could be distinguished. No DQB1 combinations met the statistical criteria for our study. In conclusion, our analysis demonstrates that mean SSM scores were not significantly different, and SSM distributions were overlapping among high- and low-risk allele combinations within loci HLA-A, B, C, DRB1, and DPB1. This analysis does not support selecting donors for HSCT recipients based on low *HistoCheck* SSM scores.

*Biol Blood Marrow Transplant 17: 1409-1415 (2011) © 2011 American Society for Blood and Marrow Transplantation*

**KEY WORDS:** HLA, Mismatched, Unrelated donor, HSCT, *HistoCheck*

## INTRODUCTION

Allogeneic hematopoietic stem cell transplantation (HSCT) is the only curative therapy for many hematologic and nonhematologic disorders. The steady expansion of unrelated stem cell donor registries has facilitated finding a matched donor for many transplant candidates, particularly those with common human leukocyte antigens (HLA) alleles and haplotypes. However, the extensive polymorphism of HLA and the remarkable disparity in the distribution of alleles and haplotypes among individuals of different ethnic and racial backgrounds remain a major hurdle for access of many patients to HSCT. A number of studies have shown that donor/recipient matching for alleles at HLA-A, -B, -C, -DRB1, and -DQB1 loci lowers the risk of clinically severe acute graft-versus-host disease (aGVHD) [1-3]. Recently, HLA-DPB1 allele mismatches were also significantly associated with an increased incidence of GVHD [4-6]. When only HLA mismatched donors are available for a given patient, the challenge becomes determining which mismatch has a less deleterious impact on clinical outcomes. Bray and colleagues [7], in a comprehensive commentary, described the National Marrow Donor Program (NMDP) guidelines for unrelated HSC donor selection including the impact of mismatches at different loci on HSCT clinical outcomes. In 2004, Elsner and colleagues [8] proposed the *HistoCheck* Web-based tool to estimate the allogenicity of mismatches with a sequence similarity matching (SSM) concept. In this concept, an SSM score (ie, allogenicity index) is generated by rating the amino acid (AA) differences between HLA allelic products based on the position within the HLA molecule and the functional similarity of AA within proteins [9]. A high SSM score (also referred to as Dissimilarity Score [DSS]) represents high dis-

similarity between HLA alleles resulting in a potentially greater deleterious impact on clinical outcomes. However, this algorithm has been challenged by 2 single-center analyses that could not associate higher SSM scores with aGVHD (in 26 patients) or in vitro T cell reactivity (in 74 patients) [10,11]. In the present study, we investigated the potential of SSM scores to predict high-risk HLA allele mismatch combinations responsible for severe aGVHD (grades III and IV) observed in a large cohort (5120 consecutive patients) of HSCT donor/recipient pairs. These allele combinations were observed in HSC transplants facilitated by the Japan Marrow Donor Program (JMDP) and published by Kawase et al. [12]. This investigation was conducted by comparing SSM scores in high-risk and low-risk allele combinations at HLA-A, -B, -C, -DRB1, and -DPB1 loci. No high-risk allele combinations at DQB1 locus met the predetermined level of statistical significance ( $P < .005$ ) and thus SSM predictions were not evaluated in this study.

## METHODS

### Identification of High- and Low-Risk HLA Allele Mismatch Combinations

Significant high-risk HLA allele mismatch combinations were identified by retrospective analysis of 5210 consecutive registered patients who underwent transplantation through the JMDP. Patient characteristics, HLA matching and typing methods, and transplant procedures are described elsewhere [12]. Briefly, 15 mismatch allele combinations were identified as high-risk allele mismatch combinations (4 at HLA-A, 1 at HLA-B, 6 at HLA-C, 1 at HLA-DRB1, and 2 at HLA-DPB1 loci). Only 1-allele mismatched pairs in the same HLA locus were considered, and

adjusted by HLA locus matching in the other loci. These mismatch combinations were found to be associated with high risk of severe aGVHD in a multivariable Cox regression model constructed with mismatch combinations and potential confounders. Confounders considered were sex, patient age, donor age, type of disease, risk of leukemia relapse, GVHD prophylaxis, and preconditioning. Each HLA mismatch combination was evaluated for each locus separately, for example, in the A\*02:06-A\*02:01 allele mismatch combination, the donor has HLA-A\*02:06, recipient has HLA-A\*02:01, and the other HLA-A allele of each donor and recipient was identical. This mismatch was compared with the HLA-A allele match. An allele mismatch combination was designated as a significant high-risk combination for severe aGVHD based on *P* values for hazard ratios (HR) for developing severe aGVHD of  $<.005$ . For example, the above mismatch combination was observed in 131 donor/recipient pairs and was associated with increased hazard of severe aGVHD (HR: 1.78, 95% confidence interval [CI]: 1.32-2.41,  $P < .001$ ). Therefore, this was considered a high-risk mismatch allele combination. On the other hand, allele mismatch combinations with a 95% CI of the HR including 1.00 were considered low-risk combinations. For example, the combination A\*24:02-A\*24:20 was observed in 60 donor/recipient pairs and was not associated with increased risk of severe aGVHD (HR: 0.64, 95% CI: 0.32-1.30,  $P = .225$ ). In addition, high-risk allele combinations observed in the context of 2 loci-linked mismatches such as (DRB1\*14:03-DQB1\*03:01) – (DRB1\*14:01-DQB1\*05:02) were excluded from this analysis because there are no explicit provisions regarding the utility of the *HistoCheck* algorithm in this setting.

### SSM Score Calculation and Comparisons

SSM scores were calculated using the *HistoCheck* tool available online at <http://www.histocheck.org/according> to the instructions posted on that Website. The averages and the distribution of SSM values were compared among high- and low-risk allele mismatch combinations in the same locus.

### Simulation Scenarios

Hypothetical scenarios were created where the recipient's HLA allele was involved in multiple allele mismatch combinations with at least 1 high-risk mismatch and 1 low-risk mismatch. These scenarios were created to simulate clinical scenarios where multiple mismatched donors are considered for a given recipient in the absence of a matched donor. In these instances, SSM scores were compared among donors from both types of mismatch combinations. In addition, to investigate the impact of the direction of the allele mismatch, we assessed whether any of the

identified 15 high-risk mismatch allele combinations associated with severe aGVHD were also a high-risk combination when considering the reverse direction of the mismatch between donor and recipient. For example, in the high-risk mismatch combination HLA-A\*26:01/26:02, the recipient had the HLA-A\*26:02 allele and the donor had the HLA-A\*26:01. In this instance, we reviewed the list of the high- and low-risk mismatch allele combinations from the Kawase study to assess the risk associated with the reverse combination (ie, recipient is HLA-A\*26:01 and donor is HLA-A\*26:02). The same assessment was performed for all identified high-risk combinations.

### Statistical Analysis

For the purpose of this analysis, low-risk mismatch allele combinations were used as a control group for SSM comparisons. SSM means and distributions were compared among aggregates of low-risk mismatch allele combinations and aggregates of high-risk combinations as well as individual high-risk combinations.


## RESULTS

### High- versus Low-Risk Allele Mismatch Combinations

In the HLA-A locus, the 4 high-risk allele mismatch combinations (observed 214 HSCT donor-recipient pairs) had a mean SSM of 2.39 (range: 1.04-4.30) compared to a mean SSM of 2.90 (range: 1.04-5.66) in 11 low-risk combinations (observed in 389 pairs). Individual high-risk combinations had the following SSM values: 1.04 (A\*02:06-02:01), 2.87 (A\*02:06-02:07), 1.36 (A\*26:02-26:01), and 4.3 (A\*26:03-26:01).

The 1 high-risk combination (B\*15:01-15:07, observed in 19 pairs) at HLA-B locus had an SSM value of 1.06. The mean SSM for 4 B low-risk combinations was 2.53 (range: 1.74-2.81). In the HLA-C locus, the mean SSM for the 7 high-risk combinations (observed in 316 pairs) was 16.60 (range: 12.36-23.86) compared to a mean SSM of 14.99 (range 1.52-23.86) in 18 low-risk combinations (observed in 578 pairs). The 1 high-risk combination (DRB1\*04:05-04:03, observed in 53 pairs) at HLA-DRB1 locus had an SSM value of 4.02. The mean SSM for 4 DRB1 low-risk combinations was 3.81 (range: 1.30-10.41). Finally, at the HLA-DPB1 locus, the mean SSM for the 2 high-risk combinations (observed in 120 pairs) was 7.47 (range: 6.98-7.95) compared to a mean SSM of 6.94 (range 1.21-12.87) in 36 low-risk combinations (observed in 1594 pairs).

A graphical representation of overall comparisons of SSM means and distributions among aggregates of low-risk mismatch allele combinations and aggregates of high-risk combinations as well as individual high-risk combinations across all loci is depicted in Figure 1. Without exception, in all these loci, the SSM



**Figure 1.** Comparison of SSM in high- versus low-risk mismatch allele combinations. SSM means are not significantly different, and SSM distributions are overlapping among high- and low-risk allele combinations within loci HLA-A, -B, -C, -DRB1, and -DPB1.

score for each high-risk combination fell within or below the range for the low-risk combinations. Two of 15 high-risk combinations had SSM scores that were lower than or equal to the lowest SSM score in any of the low-risk combinations in the corresponding locus. A detailed list of individual low-risk combinations is presented in the original publication by Kawase et al. [12].

### Simulation Scenarios

To simulate clinical scenarios where multiple mismatched donors are considered for a given recipient, SSM values for high-risk and low-risk combinations including the same recipient allele were compared. The recipient allele in 13 of the 15 identified high-risk mismatch allele combinations (Loci HLA-A\*, C\*, DRB1\*, and DPB1\*) was included in 1 or more low-risk allele mismatch combinations (Table 1). In 1 scenario, a recipient with an A\*02:07 allele had 2 potential mismatched donors at this allele for either A\*02:06 (high-risk mismatch allele combination) or A\*02:01 (low-risk mismatch allele combination). SSM values for the high-risk and the low-risk combinations were 2.90 and 1.83, respectively, as one might expect. However, in another scenario, a recipient with an A\*02:01 allele had 2 potential mismatched donors at this allele for either A\*02:06 (high-risk mismatch allele combination) or A\*02:07 (low-risk mismatch allele combination), the opposite was observed. SSM values for the high- and the low-risk combinations were 1.04 and 1.83, respectively. All simulation scenarios are summarized in Table 1.

Overall, no predictable SSM pattern of association with high- or low-risk mismatch allele combinations could be distinguished. In all but 2 instances (at HLA-A\* and -DRB1\* loci), at least 1 low-risk combination scored a higher SSM value than the high-risk

mismatch combination including the same recipient allele.

### Direction of the Allele Mismatch Combinations

None of the 15 high-risk combinations were high risk in both directions. In 10 of these combinations, when the direction of the allele mismatch was reversed between donor and recipient, there was no longer an association with severe aGVHD (Table 2). An example of such a unidirectional risk is the combination where the recipient has A\*02:01 and the donor has A\*02:06 (high risk) versus the recipient having A\*02:06 and the donor having A\*02:01 (low risk). Both of these combinations have an SSM value of 1.04 according to *HistoCheck*.

### DISCUSSION

Optimal effectiveness and safety of HSCT requires high degree of HLA allele matching between donors and recipients [7,13]. However, many patients who need HSCT do not have an HLA-matched donor, which lowers the probability of cure. Undoubtedly, a predictive algorithm for definition of “low-risk” HLA mismatches has the potential of broadening the use of mismatched donors and increasing the availability of unrelated donors. Several approaches have been proposed to identify such low-risk mismatches, with the simplest being a comparison between alleles based on the number of amino acid mismatches. However, prior reports have not found any evidence for selecting an allele with a lower number of AA substitutions of its allelic product [14,15]. Although hypothetically appealing, the notion that the lower the number of AA substitutions the more low-risk the mismatch is



**Table 1. Simulation Scenarios for Comparison between SSM Scores among Multiple High- versus Low-Risk Mismatched Donors Potentially Available for a Patient with a Given HLA Allele**

HLA Locus	MM Combination (Donor-Patient)	N	HR (95% CI)	P	Risk Group	SSM
A	02:06-02:01	131	1.78 (1.32-2.41)	<.001	High	1.04
	02:07-02:01	20	1.12 (0.42-3.02)	.81	Low	1.83
	02:06-02:07	27	3.45 (2.09-5.70)	<.001	High	2.90*
	02:01-02:07	28	0.83 (0.34-2.03)	.70	Low	1.83
C	04:01-03:03	42	2.81 (1.72-4.60)	.001	High	23.86
	08:01-03:03	80	2.32 (1.58-3.40)	.001	High	16.65
	07:02-03:03	18	2.16 (0.96-4.85)	.06	Low	21.90
	03:04-03:03	62	0.83 (0.41-1.68)	.61	Low	1.52
	03:04-08:01	69	2.34 (1.55-3.52)	.001	High	7.13
	03:03-08:01	76	1.07 (0.63-1.84)	.78	Low	16.65
	14:02-03:04	23	3.66 (2.00-6.68)	.001	High	17.30
	15:02-03:04	27	3.77 (2.20-6.47)	.001	High	12.36
	08:01-03:04	47	1.64 (0.98-2.76)	.06	Low	15.13
	01:02-03:04	12	1.85 (0.59-5.81)	.29	Low	18.30
	07:02-03:04	33	1.22 (0.58-2.59)	.59	Low	20.38
	03:03-03:04	83	1.08 (0.63-1.85)	.76	Low	1.52
	03:03-15:02	25	3.22 (1.75-5.89)	.001	High	13.88
	08:01-15:02	36	1.59 (0.79-3.21)	.19	Low	15.33
	04:05-04:03	53	2.13 (1.28-3.53)	.003	High	4.02*
	04:10-04:03	17	1.01 (0.32-3.21)	.98	Low	2.50
	04:06-04:03	30	0.99 (0.46-2.10)	.99	Low	1.45
	05:01-09:01	71	2.03 (1.30-3.16)	.002	High	7.95
DRB1	04:02-09:01	17	0.33 (0.04-2.36)	.27	Low	11.19
	02:01-09:01	47	1.37 (0.75-2.51)	.30	Low	9.98
DPB1	03:01-09:01	15	0.8 (0.19-3.22)	.75	Low	3.98
	04:01-09:01	11	0.9 (0.22-3.66)	.89	Low	12.87
	03:01-05:01	49	2.41 (1.49-3.89)	<.001	High	6.98
	06:01-05:01	13	2.5 (0.92-6.77)	.07	Low	7.95
	04:02-05:01	79	1.47 (0.90-2.40)	.12	Low	6.70
	02:02-05:01	41	0.43 (0.13-1.35)	.15	Low	6.18
	09:01-05:01	48	0.71 (0.29-1.73)	.46	Low	7.95
	04:01-05:01	29	0.73 (0.23-2.29)	.59	Low	5.46
	14:01-05:01	26	1.17 (0.48-2.84)	.73	Low	7.77

N indicates number of donor-patient pairs in whom the mismatch allele combination was observed; HR, hazard ratio of developing severe acute graft-versus-host disease (aGVHD) compared to matched pairs as described in Kawase et al. [12]; SSM, sequence similarity matching; CI, confidence interval.

P values: for the corresponding estimated hazard ratio.

In all but 2 instances (marked with \*), at least 1 low-risk combination scored a higher SSM value than the high-risk mismatch combination including the same recipient allele.

\*The only 2 instances where SSM value was highest for the combination associated severe aGVHD.

challenged by 2 lines of evidence. First, a recent NMDP registry analysis suggested no difference in outcome when comparing antigen mismatch to allele level (high resolution) mismatch within the same antigen group [13]. Second, at least 2 reports have indicated that severe aGVHD can occur in the presence of only 1 AA mismatch between the donor and recipient allelic products [12,16].

In addition, at least 2 “epitope” based approaches have been proposed. One is based on serologically crossreactive groups (CREG) of antigens, and the other is based on comparisons of “functional epitope” structure through molecular viewing of the HLA structure and the determination of antibody-accessible polymorphic AAs (HLAMatchmaker) [17-19]. Neither of these 2 approaches have yielded predictions that were associated with a survival benefit in patients who underwent mismatched hematopoietic cell transplants from unrelated donors in registry analyses [20,21].

In the current analysis, we investigated whether *HistoCheck* SSM scores can predict 15 high-risk HLA

allele mismatch combinations responsible for severe aGVHD observed in 5120 consecutive HSCT donor-recipient pairs facilitated through JMDP. It is noteworthy that the association between these 15 allele combinations and high risk for severe aGVHD has not yet been validated in an independent patient population; however, we believe that this association is reasonably robust because of the conservative approach and the rigorous statistical methods pursued in assigning these associations as described elsewhere [12]. SSM score comparisons were performed between high-risk and low-risk combinations at loci HLA-A, -B, -C, -DRB1, and -DPB1. Significant overlap exists between the high- and low-risk mismatches with respect to SSM scores. In all investigated loci, the SSM score for each high-risk combination fell within or below the range for the low-risk combinations, thus demonstrating the unreliability of SSM scores with respect to aGVHD risk. In addition, in 2 high-risk combinations (13%), SSM scores were less than or equal to the lowest SSM score in any of the low-risk combinations in the

**Table 2. Mismatch Allele Combinations Where the Risk of aGVHD Depends on the Direction of the Mismatch**

HLA Locus	MM Combination (Donor-Patient)	N	HR (95% CI)	P	Risk Group	SSM
A*	02:06-02:01	131	1.78 (1.32-2.41)	<.001	High	1.04
	02:01-02:06	138	1.23 (0.87-1.73)	.223	Low	
	02:06-02:07	27	3.45 (2.09-5.70)	<.001	High	2.87
	02:07-02:06	22	0.71 (0.23-2.24)	.571	Low	
	26:02-26:01	21	3.35 (1.89-5.91)	<.001	High	1.36
	26:01-26:02	24	0.64 (0.26-1.58)	.34	Low	
	26:03-26:01	35	2.17 (1.29-3.64)	.003	High	4.30
	26:01-26:03	34	1.37 (0.73-2.57)	.326	Low	
	08:01-03:03	80	2.32 (1.58-3.40)	.001	High	16.65
	03:03-08:01	76	1.07 (0.63-1.84)	.782	Low	
C*	03:04-08:01	69	2.34 (1.55-3.52)	.001	High	15.13
	08:01-03:04	47	1.64 (0.98-2.76)	.057	Low	
	04:01-03:03	42	2.81 (1.72-4.60)	.001	High	23.86
	03:03-04:01	31	1.73 (0.89-3.36)	.103	Low	
	04:05-04:03	53	2.13 (1.28-3.53)	.003	High	4.02
DRB1*	04:03-04:05	54	1.27 (0.74-2.20)	.379	Low	
DPB1*	05:01-09:01	71	2.03 (1.30-3.16)	.002	High	7.95
	09:01-05:01	48	0.71 (0.29-1.73)	.457	Low	
	03:01-05:01	49	2.41 (1.49-3.89)	<.001	High	6.98
	05:01-03:01	83	1.20 (0.75-1.94)	.434	Low	

N indicates number of donor-patient pairs in whom the mismatch allele combination was observed; HR, hazard ratio of developing severe acute graft-versus-host disease (aGVHD) compared to matched pairs as described in Kawase et al. [12]; SSM, sequence similarity matching; CI, confidence interval.

P values: for the corresponding estimated hazard ratio.

None of the 15 high-risk combinations were high risk in both directions. In 10 of these combinations, when the direction of the allele mismatch was reversed between donor and recipient, there was no longer an association with severe aGVHD.

\*The only 2 instances where SSM value was highest for the combination associated with severe aGVHD.

corresponding locus. Furthermore, when simulation scenarios were constructed for recipients included in both high- and low-risk combinations, one would expect that high-risk mismatch combinations have higher SSM values compared to low-risk combinations of the same recipient allele. However, only 2 of the 15 high-risk combinations (13%) had higher SSM scores than alternative low-risk combinations potentially available for a patient with alleles included in these combinations. Interestingly, none of the identified 15 mismatch allele combinations were associated with high-risk for severe aGVHD in both directions. Ten of these combinations were no longer associated with severe aGVHD when the direction of the allele mismatch was reversed between the donor and the recipient. However, the SSM score remains unchanged because *HistoCheck* does not distinguish the direction of the allele mismatch between donor and recipient.

The lack of association between SSM and clinical data in our study may be attributed at least in part to the limitations acknowledged by the developers of the algorithm. Namely, underestimating the impact of single AA mismatches, not accounting for clinical data suggesting disproportionately larger impact of substitutions at given positions, particularly position 116, and assignment of positions based on crystallographic data of HLA-A2 for class I and of HLA-DR1 for class II alleles [8]. In addition, *HistoCheck* by design does not account for the direction of the allele mismatch, which has shown to be clinically significant [12].

Limitations of our study include relatively few numbers of high-risk allele mismatch combinations,

not including mismatch combinations at the HLA-DQB1 locus, the potential for misclassification of some mismatch combinations as low risk because of relatively small number of subjects in subcategories, and inclusion of patients only from 1 ethnic background. Nevertheless, these results and previous reports do not support the utilization of *HistoCheck* predictions in unrelated donor selection [10,11].

In conclusion, safely maximizing access to mismatched HSCT unrelated donors requires a robust understanding of the rules that govern permissible HLA mismatching, and a better understanding of HLA-associated risks of GVHD. The lack of association between the *HistoCheck* predictions and all previously proposed prediction algorithms and the observed clinical outcomes strongly emphasizes the utmost importance of clinical validation of any prediction algorithm prior to its utilization in clinical patient care. Moving forward, it is prudent in evaluating any prediction algorithm to rely primarily on clinical correlations rather than simply putative biological plausibility.

## ACKNOWLEDGMENTS

The authors acknowledge the fact that this study could not have been conducted but for the original findings of the study by Dr. Takakazu Kawase and his colleagues, and are grateful for their entire research group involved in the original study. The authors also thank Dr. Brian Bolwell, Chairman of Taussig Cancer Institute at Cleveland Clinic, for his insightful revision

of the manuscript. The authors report no conflicts of interest for this study.

**Financial disclosure:** The authors have nothing to disclose.

## REFERENCES

- Petersdorf EW, Anasetti C, Martin PJ, et al. Limits of HLA mismatching in unrelated hematopoietic cell transplantation. *Blood*. 2004;104:2976-2980.
- Morishima Y, Sasazuki T, Inoko H, et al. The clinical significance of human leukocyte antigen (HLA) allele compatibility in patients receiving a marrow transplant from serologically HLA-A, HLA-B, and HLA-DR matched unrelated donors. *Blood*. 2002;99:4200-4206.
- Flomenberg N, Baxter-Lowe LA, Confer D, et al. Impact of HLA class I and class II high-resolution matching on outcomes of unrelated donor bone marrow transplantation: HLA-C mismatching is associated with a strong adverse effect on transplantation outcome. *Blood*. 2004;104:1923-1930.
- Ludajic K, Balavarca Y, Bickeboller H, et al. Impact of HLA-DPB1 allelic and single amino acid mismatches on HSCT. *Br J Haematol*. 2008;142:436-443.
- Kawase T, Matsuo K, Kashiwase K, et al. HLA mismatch combinations associated with decreased risk of relapse: implications for the molecular mechanism. *Blood*. 2009;113:2851-2858.
- Crocchiolo R, Zino E, Vago L, et al. Nonpermissive HLA-DPB1 disparity is a significant independent risk factor for mortality after unrelated hematopoietic stem cell transplantation. *Blood*. 2009;114:1437-1444.
- Bray RA, Hurley CK, Kamani NR, et al. National marrow donor program HLA matching guidelines for unrelated adult donor hematopoietic cell transplants. *Biol Blood Marrow Transplant*. 2008;14:45-53.
- Elsner HA, DeLuca D, Strub J, Blasczyk R. HistoCheck: rating of HLA class I and II mismatches by an internet-based software tool. *Bone Marrow Transplant*. 2004;33:165-169.
- Elsner HA, Blasczyk R. Sequence similarity matching: proposal of a structure-based rating system for bone marrow transplantation. *Eur J Immunogenet*. 2002;29:229-236.
- Shaw BE, Barber LD, Madrigal JA, Cleaver S, Marsh SG. Scoring for HLA matching? A clinical test of HistoCheck. *Bone Marrow Transplant*. 2004;34:367-368, author reply 369.
- Heemskerk MB, Doxiadis II, Roelen DL, Claas FH, Oudshoorn M. The HistoCheck algorithm does not predict T-cell alloreactivity in vitro. *Bone Marrow Transplant*. 2005;36:927-928.
- Kawase T, Morishima Y, Matsuo K, et al. High-risk HLA allele mismatch combinations responsible for severe acute graft-versus-host disease and implication for its molecular mechanism. *Blood*. 2007;110:2235-2241.
- Lee SJ, Klein J, Haagenson M, et al. High-resolution donor-recipient HLA matching contributes to the success of unrelated donor marrow transplantation. *Blood*. 2007;110:4576-4583.
- Heemskerk MB, Roelen DL, Dankers MK, et al. Allo-genetic MHC class I molecules with numerous sequence differences do not elicit a CTL response. *Hum Immunol*. 2005;66:969-976.
- Heemskerk MB, Cornelissen JJ, Roelen DL, et al. Highly diverged MHC class I mismatches are acceptable for haematopoietic stem cell transplantation. *Bone Marrow Transplant*. 2007;40:193-200.
- Morishima Y, Kawase T, Malkki M, Petersdorf EW. Effect of HLA-A2 allele disparity on clinical outcome in hematopoietic cell transplantation from unrelated donors. *Tissue Antigens*. 2007;69(Suppl 1):31-35.
- Rodey GE, Fuller TC. Public epitopes and the antigenic structure of the HLA molecules. *Crit Rev Immunol*. 1987;7:229-267.
- Duquesnoy RJ, Marrari M. HLAMatchmaker: a molecularly based algorithm for histocompatibility determination. II. Verification of the algorithm and determination of the relative immunogenicity of amino acid triplet-defined epitopes. *Hum Immunol*. 2002;63:353-363.
- Duquesnoy RJ, Askar M. HLAMatchmaker: a molecularly based algorithm for histocompatibility determination. V. Eplet matching for HLA-DR, HLA-DQ, and HLA-DP. *Hum Immunol*. 2007;68:12-25.
- Wade JA, Hurley CK, Takemoto SK, et al. HLA mismatching within or outside of cross-reactive groups (CREGs) is associated with similar outcomes after unrelated hematopoietic stem cell transplantation. *Blood*. 2007;109:4064-4070.
- Duquesnoy R, Spellman S, Haagenson M, Wang T, Horowitz MM, Oudshoorn M. HLAMatchmaker-defined triplet matching is not associated with better survival rates of patients with class I HLA allele mismatched hematopoietic cell transplants from unrelated donors. *Biol Blood Marrow Transplant*. 2008;14:1064-1071.